

Mitochondrial reactive oxygen species and Ca²⁺ signaling

Cristina Camello-Almaraz, Pedro J. Gomez-Pinilla, Maria J. Pozo, and Pedro J. Camello

Department of Physiology, Faculty of Veterinary Sciences and Nursing School, University of Extremadura, Cáceres, Spain

Submitted 29 April 2006; accepted in final form 1 June 2006

Camello-Almaraz, Cristina, Pedro J. Gomez-Pinilla, Maria J. Pozo, and Pedro J. Camello. Mitochondrial reactive oxygen species and Ca²⁺ signaling. *Am J Physiol Cell Physiol* 291: C1082–C1088, 2006; doi:10.1152/ajpcell.00217.2006.—Mitochondria are an important source of reactive oxygen species (ROS) formed as a side product of oxidative phosphorylation. The main sites of oxidant production are complex I and complex III, where electrons flowing from reduced substrates are occasionally transferred to oxygen to form superoxide anion and derived products. These highly reactive compounds have a well-known role in pathological states and in some cellular responses. However, although their link with Ca²⁺ is well studied in cell death, it has been hardly investigated in normal cytosolic calcium concentration ([Ca²⁺]_i) signals. Several Ca²⁺ transport systems are modulated by oxidation. Oxidation increases the activity of inositol 1,4,5-trisphosphate and ryanodine receptors, the main channels releasing Ca²⁺ from intracellular stores in response to cellular stimulation. On the other hand, mitochondria are known to control [Ca²⁺]_i signals by Ca²⁺ uptake and release during cytosolic calcium mobilization, specially in mitochondria situated close to Ca²⁺ release channels. Mitochondrial inhibitors modify calcium signals in numerous cell types, including oscillations evoked by physiological stimulus. Although these inhibitors reduce mitochondrial Ca²⁺ uptake, they also impair ROS production in several systems. In keeping with this effect, recent reports show that antioxidants or oxidant scavengers also inhibit physiological calcium signals. Furthermore, there is evidence that mitochondria generate ROS in response to cell stimulation, an effect suppressed by mitochondrial inhibitors that simultaneously block [Ca²⁺]_i signals. Together, the data reviewed here indicate that Ca²⁺-mobilizing stimulus generates mitochondrial ROS, which, in turn, facilitate [Ca²⁺]_i signals, a new aspect in the biology of mitochondria. Finally, the potential implications for biological modeling are discussed.

mitochondria; calcium

MITOCHONDRIA SERVE as the main source for ATP used by eukaryotic cells in the course of cellular functions. However, developments in imaging techniques have led to characterization of mitochondria involvement in other key processes, such as calcium signals or apoptosis. The number of excellent reviews and references reporting on these aspects is very high, and a fast bibliographic search will yield literally thousands of references.

A consequence of mitochondria function is the production of reactive oxygen species (ROS) during physiological and pathological states (Fig. 1). This process has also been extensively studied because ROS have a clear role in mediating functional alterations of cell physiology in inflammation, ischemia, aging, and other conditions (19, 56, 57). There is also an emerging field of study of ROS control of cellular functions (18). The present review summarizes evidence for a poorly studied

aspect of mitochondria biology, the role of ROS generated during oxidative phosphorylation in Ca²⁺ signals.

Mitochondrial Production of Reactive Oxygen Species

“Reactive oxygen species” refers to a group of oxygen-containing compounds with the ability to react with reducible compounds. They comprise superoxide (O₂⁻·), hydrogen peroxide (H₂O₂), and the highly reactive hydroxyl radical (·OH), although minor amounts of singlet oxygen can also be formed by cells. The initial product of the electron transport chain (ETC) is O₂⁻·, which is quickly transformed into H₂O₂ by the enzyme superoxide dismutase (SOD). H₂O₂ can be reduced to water by catalase or glutathione peroxidase or can be converted into ·OH in presence of reduced transition metals (reduced copper or iron).

The main source of O₂⁻· is respiratory complexes I and III located at the inner mitochondrial membrane (12, 56, 83). These complexes generate a small amount of O₂⁻· as a side product of electron transport during oxidative phosphorylation. O₂⁻· is released into the matrix in the case of complex I and to both the matrix and the intermembranous space by complex III. Complex III forms O₂⁻· during cycling of the electron acceptor ubiquinone, which can donate electrons to molecular oxygen in both the internal and the external face of the inner mitochondrial membrane.

The relative importance of these sites in O₂⁻· mitochondrial output depends on tissue and mitochondria respiratory status. For example, in fully respiring mitochondria from heart muscle (characterized by high electron flow, fast ATP synthesis, partial depolarization, and a decreased NADH-to-NAD⁺ ratio, corresponding to state 3), complex III seems to be dominant, and the generation of O₂⁻· is proportional to electron flow rate (84). In this situation oxygen, reduced substrates, and downstream electron acceptors are needed. Therefore, inhibition of electron flow into the ubiquinone cycle of complex III (by inhibition of complex I with rotenone, blockade of the cycle with myxothiazol, or inhibition of downstream electron acceptor cytochrome *c*) reduces O₂⁻· generation, whereas antimycin enhances it by building up the partially reduced form of ubiquinone.

Complex I is the main source of O₂⁻· in state 4, when electron transport rate and ATP synthesis are low and substrates are highly reduced (high NADH-to-NAD⁺ ratio). The redox potential of complex I seems to be even higher than the NADH/NAD⁺ couple, which renders it a thermodynamically unstable center prone to electron leakage to oxygen (47). In this center, O₂⁻· production is increased in any situation leading to reduction of ETC components (for example, by application of rotenone, which blocks complex I distal to the O₂⁻· produc-

Address for reprint requests and other correspondence: P. J. Camello, Dept. of Physiology, Fac. Veterinary Sci., Avda Universidad s/n, 10071 Cáceres, Spain (e-mail: pcamello@unex.es).

The costs of publication of this article were defrayed in part by the payment of page charges. The article must therefore be hereby marked “advertisement” in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.




Fig. 1. Main mechanisms for mitochondrial reactive oxygen species (ROS) generation. Reduced substrates synthesized in metabolic pathways supply electrons (e^-) to complex I and II of the electron transport chain. The main centers for superoxide ($O_2^{\cdot-}$) formation are complex I and III, although small amounts can be formed at complex II and IV (omitted for clarity). The electron carrier of complex III ubiquinone (Q) is reduced to ubiquinol (QH_2), which transfers an electron to cytochrome *c* (Cy C) through a iron protein (not shown) inhibited by myxothiazol (Myx). The resulting semiquinone (Q^*) is oxidized back to ubiquinone by cytochrome *b* (Cy b), and can also transfer electrons to oxygen to form $O_2^{\cdot-}$. Myxothiazol reduces $O_2^{\cdot-}$ production because it blocks Q^* formation, whereas antimycin A (Ant A) enhances it by increasing Q^* levels. Rotenone (Rot) inhibits electron flow distal to $O_2^{\cdot-}$ generation, enhancing its production. The main routes for $O_2^{\cdot-}$ transformation are represented. SOD, superoxide dismutase; H_2O_2 , hydrogen peroxide; $ONOO^-$, peroxynitrite; NO^* , nitric oxide; $\Delta\Psi_m$, mitochondrial potential; AS, ATP synthase. Only the inner mitochondrial membrane is represented.

tion point, or inhibition of electron flow at the level of cytochrome *c*).

Superoxide anions in the mitochondrial matrix are quickly dismutated by manganese-containing SOD, whereas those in the intermembranous space are converted by Zn- or Cu-SOD (83). If the $O_2^{\cdot-}$ concentration in the matrix is high enough, part of the anion escapes to the intermembranous space and cytosol via anion channels (3).

The above-described mechanisms are supported by experimental results from isolated mitochondria studies. Results from mitochondria in living, intact cells support a main role of complex III as ROS source in several systems, because rotenone or myxothiazol can impair ROS production in active cells (see, e.g., Refs. 2, 3, 10, 17). Together, these results suggest that mitochondria in these cells are in state 3. On the other hand, in other preparations administration of rotenone increases ROS production, probably through complex I, indicative of mitochondria in state 4. These opposite results can be explained by mitochondria switching between states 3 and 4 (83). In fact, it has been shown that heart mitochondria display oscillatory behavior, with cycles in mitochondrial potential and production of ROS (2, 3).

Cellular stimulation leads to fast increases in reduced substrates derived from the Krebs cycle (NADH and $FADH_2$) in the mitochondrial matrix, which has been related to cytosolic calcium concentration ($[Ca^{2+}]_i$) increases within mitochondria (28, 31, 76, 86). At the same time this situation will favor increased electron flow and rate of ATP demand (used for Ca^{2+} homeostasis, contraction, exocytosis, etc.), which would facilitate ROS production at complex III (15). Several systems have been reported to increase ROS generation in response to a variety of stimuli (10, 33, 38, 42, 43, 65, 88, 89, 91).

In any case, it has been argued that there is no convincing mechanism to explain how calcium signals can enhance $O_2^{\cdot-}$ generation, given that a rise in mitochondrial calcium concentration depolarizes mitochondria and should decrease $O_2^{\cdot-}$ formation (55, 77, 93), similar to the effect of uncouplers (24, 56, 91). Alternative proposals for Ca^{2+} -evoked increases in ROS production are direct control of ETC complexes or inhibition of complex I and IV by nitric oxide and peroxy-

trite, which can be generated by mitochondria (6). However, the available results on this issue are conflicting, because some reports describe ROS generation in response to mitochondrial uncouplers (38) whereas others show that mitochondrial Ca^{2+} accumulation decreases it (78). Overall, the data show that calcium induces ROS production if mitochondria are treated with some inhibitor but reduces it under normal conditions (6). A possible explanation for contradictory results would be that mitochondria in vivo are under some kind of tonic inhibitory influence, allowing imported calcium to enhance $O_2^{\cdot-}$ formation in certain conditions (see also *Mitochondrial Production of Reactive Species Participates in Normal Ca^{2+} Signals*).

Mitochondria Participate in Ca^{2+} Homeostasis

Participation of mitochondria in calcium signal has been revealed by two main experimental approaches: 1) mitochondria uptake and release of Ca^{2+} in the course of Ca^{2+} mobilization and 2) inhibition of mitochondrial function modifying the shape of Ca^{2+} signals. Both fluorescent and chemiluminescent methods, including targeted optical sensors and confocal microscopy, have allowed a detailed characterization of mitochondrial calcium mobilization (71). Regarding manipulation of mitochondrial function, most studies rely on the use of specific inhibitors of the ETC or chemical uncouplers of the mitochondrial potential. A number of authoritative reviews are available (see, for example Refs. 20, 30, 55, 66, 68, 71).

Mitochondria behave as a high-capacity, low-affinity transient calcium store. Changes in $[Ca^{2+}]_i$ in response to hormones and neurotransmitters induce increases in the mitochondrial matrix calcium concentration ($[Ca^{2+}]_m$). Calcium ions enter across the inner mitochondrial membrane following its electrochemical gradient when $[Ca^{2+}]_i$ levels reach a submicromolar threshold (55, 76). The route is called the Ca^{2+} uniporter, an elusive pathway that remains poorly understood, although a recent report identifies it as a calcium-selective channel (44). In addition, it has been reported that Ca^{2+} can enter mitochondria through a "rapid mode" mechanism (75), on a millisecond timescale, as well as through a mitochondrial isoform of ryanodine receptor described in some excitable cells (4). Both sustained and transient (even oscillatory) $[Ca^{2+}]_m$

responses have been reported in multiple cell types (9, 20, 31, 70). The Ca^{2+} accumulated in mitochondria during the rising phase of the $[\text{Ca}^{2+}]_i$ signal is subsequently released during the declining phase of the signals via a $\text{Na}^+/\text{Ca}^{2+}$ exchanger (54, 72), although both the Ca^{2+} uniporter and the permeability transition pore have also been proposed as efflux routes (29, 54, 59).

In essence, this pattern of $[\text{Ca}^{2+}]_m$ changes is considered to be due to the close apposition between mitochondria and calcium release channels of endoplasmic reticulum (68, 69) [inositol 1,4,5-trisphosphate (IP_3) and ryanodine receptors (IP_3R and RyR)], as well as proximity between mitochondria and plasma membrane (Fig. 2). This arrangement allows mitochondria to take up calcium ions entering the cytosol from internal stores or even from extracellular space. This implies that mitochondria partially buffer Ca^{2+} transients but also participate in spatiotemporal propagation of signals, because they modulate the Ca^{2+} microdomains in the immediate vicinity of calcium channels, which are known to be sensitive to calcium concentration. Thus mitochondria operate either as a barrier buffer (9, 11, 32, 60, 79, 82) or as a facilitating factor in the spreading of calcium signals (20, 73, 74), behaving as an effective shaper of calcium signals elicited by IP_3R and RyR activated during neurohormonal challenge. Although some authors report that this mechanism is also true for capacitative calcium influx in some models (32), we have shown (9) that in




Fig. 2. Interaction between mitochondria and Ca^{2+} -release channels. After cell stimulation, inositol 1,4,5-trisphosphate (IP_3) and cADP ribose (cADPr) release calcium from endoplasmic reticulum (ER) through their respective receptors [IP_3R and ryanodine receptors (RyR)]. Local cytosolic calcium concentration ($[\text{Ca}^{2+}]_i$) increases are buffered by mitochondrial uptake through the calcium uniporter (MU) and possible alternative routes (not shown). Intramitochondrial calcium activates synthesis of reduced substrates (NADH) by metabolic pathways and accelerates the electron transport chain, increasing ROS production, which in turn facilitates Ca^{2+} release by sensitization of IP_3R and RyR . Other putative mechanisms increasing ROS generation are modification of electron transport complexes by mitochondrial calcium and inhibition by nitric oxide (NO^*), which can be formed by Ca^{2+} and by mitochondria. Mitochondria can also support Ca^{2+} release by efflux via $\text{Na}^+/\text{Ca}^{2+}$ exchanger (NCX) and by increase of cADP ribose. Outer mitochondrial membrane is not represented.

exocrine cells is a wrong conclusion, based on the misleading effect of mitochondrial buffer.

Mitochondrial calcium increases also have consequences for mitochondria, mainly acceleration of metabolic enzymes (see review in Ref. 6) and activation of the permeability transition pore, a mechanism whereby permeability of the inner membrane increases dramatically in response to factors such as oxidative stress or calcium overload. The former response increases production of reduced substrates for ETC-mediated ATP synthesis, whereas permeability transition releases not only calcium but also high-molecular-weight compounds, such as cytochrome *c* and other proteins, which in turn activates apoptotic signals (6). Some reports have described fast and reversible activation of permeability transition and have proposed a physiological role for this process (36, 59, 85).

Ca²⁺ Homeostasis Components Are Modified by Reactive Oxygen Species

Calcium transport systems are sensitive to redox conditions. Thus oxidants inhibit Ca^{2+} extrusion via plasma membrane Ca^{2+} -ATPase (PMCA) (48, 92) and Ca^{2+} reuptake into sarco(endo)plasmic reticulum Ca^{2+} -ATPase (SERCA)-operated stores (41, 49), whereas there are conflicting reports about the effects in the plasma membrane $\text{Na}^+/\text{Ca}^{2+}$ exchange (35, 46).

Superoxide and other ROS inhibit L-type calcium channels (see, e.g., Ref. 26), acting directly on the channel, probably on the voltage sensor domain, although this conclusion is based on indirect data (46). An indirect mechanism of action of ROS on voltage-operated calcium channels through modifications of plasma membrane potential cannot be ruled out, as electrophysiological data show ROS-induced changes in potassium currents that would modify plasma membrane potential (reviewed in Ref. 46).

Regarding the main channels for calcium release from internal stores, thiol oxidants acting on sulfhydryl residues release calcium through IP_3R (5, 7, 52) and RyR (1, 23, 80), a mechanism that enhances (and even elicits) Ca^{2+} oscillations in exocrine cells (81, 90) and HeLa cells (5). This process could be due to sensitization of IP_3R -mediated calcium release (51, 52) to resting IP_3 levels, a phenomenon that has been integrated in a model to facilitate calcium oscillations (52). This effect of oxidizing reagents has been experimentally observed at the single-channel level for RyR (46). Together, the available evidence shows that the presence of sulfhydryl groups in these receptor channels enables them to respond to low levels of oxidants far below those achieved during pathological states or in extreme experimental conditions. Another example of facilitation by oxidants is the cooperative effect of H_2O_2 on the activation of TRPM2 Ca^{2+} -permeable channels by ADP ribose (45).

As a whole, the available evidence shows that ROS can induce $[\text{Ca}^{2+}]_i$ increases by release from internal stores and impairment of Ca^{2+} clearance systems. In fact, $[\text{Ca}^{2+}]_i$ increase is a constant feature of pathological states associated with oxidative stress.

Mitochondrial Production of Reactive Species Participates in Normal Ca²⁺ Signals

The relationship between oxidative stress and calcium signals has been extensively studied, given the pathological rele-

vance of the field. However, although the concept of endogenous ROS as intracellular messengers is not recent (18, 53), their involvement in physiological calcium signals has been only marginally addressed.

Physiological levels of stimulation generate in several systems transient $[Ca^{2+}]_i$ oscillations, as is the case for the pancreatic acinar cells in response to postprandial concentration of the gut hormone CCK (63). Other examples of physiological $[Ca^{2+}]_i$ signals are fast and localized, nonpropagating signals observed in excitable (Ca^{2+} sparks; Refs. 39, 64) and nonexcitable (Ca^{2+} puffs; Ref. 11) cells. These subcellular calcium signals play clear regulatory roles in several systems (e.g., control of sarcolemma potential in muscle cells or exocytosis in exocrine pancreas). Participation of mitochondria in calcium oscillations has been evidenced by the inhibitory effect of mitochondrial inhibitors and uncouplers in pancreatic acinar cells stimulated by CCK (9, 79) and gonadotropes activated by gonadotropin-releasing hormone (40). In the same line is the finding that some antihypertensive potassium channel openers activate $[Ca^{2+}]_i$ sparks in artery through mitochondria-formed ROS (91). Another recent article reports suppression of serum-induced $[Ca^{2+}]_i$ oscillations in bone marrow cells by mitochondrial depolarization (25). Following on this line of evidence, we proposed (8) a model in which mitochondrial oxidants play a cooperative role in hormonally induced $[Ca^{2+}]_i$ oscillations in pancreatic acinar cells.

To establish a role for mitochondrial ROS it is necessary, however, to demonstrate that 1) blockade or scavenging of ROS modifies the calcium signal and 2) ROS production takes place during physiological stimulus. Both types of results are already available.

If the model supported by these results is true, disruption of the ability of mitochondria to produce small amounts of oxidants would alter normal Ca^{2+} signals: too much ROS production would induce massive and sustained Ca^{2+} release, whereas loss of ROS production would deprive Ca^{2+} release channels of an endogenous sensitizer. In recent years several groups have reported that antioxidants or interference with mitochondrial ROS production modifies calcium signals. We have recently shown in pancreatic acinar cells that a mitochondria-targeted antioxidant (10) and mitochondrial inhibitors (9, 10) reduce CCK-evoked Ca^{2+} oscillation and ROS formation (10). A similar mechanistic evidence was provided in smooth muscle, where ROS production and Ca^{2+} sparks induced by diazoxide are blocked by rotenone and FCCP (91). Additional lines of evidence supporting this role of mitochondria are findings that pretreatment with *N*-acetylcysteine inhibits CCK-evoked oscillations (8), antioxidants reduce spark frequency in skeletal muscle (37), and $[Ca^{2+}]_i$ signals are evoked by hypoxia in pulmonary artery (88). Although not directly related to mitochondria, other reports also reinforce this concept: blockade of extramitochondrial ROS suppresses histamine-evoked $[Ca^{2+}]_i$ oscillations in endothelial cells (34), and there is a clear correlation between ROS production and spreading of calcium waves in alga cells (13). Mitochondrial blockers also prevent nongenomic effects of estrogen such as $[Ca^{2+}]_m$ increase (61) and cell proliferation (21).

The ability of cellular agonists to enhance mitochondrial ROS has been reported in several systems. We have shown (10) that postprandial levels of the gut hormone CCK increase ROS in the mitochondria of intact pancreatic acinar cells, an

effect blocked by rotenone. Mitochondrial production of ROS has also been reported in cultured cells and endothelial cells for estrogen (21, 22, 61), in pulmonary arteries under hypoxia (88, 89), and in cardiac and vascular muscles and endothelial cells for angiotensin II (42, 43, 65). Similar results have been reported for hyperglycemia in rat kidney tubular cells (33) and tumor necrosis factor in carcinoma cells (38). As mentioned above, some vasodilators generate mitochondrial ROS to elicit $[Ca^{2+}]_i$ sparks in artery myocytes (91).

Although most of the evidence described above is rather correlative, some reports provide a mechanistic approach (10, 91). Together with the facilitation of Ca^{2+} release by exogenous oxidants (see *Ca²⁺ Homeostasis Components Are Modified by Reactive Oxygen Species*), the described evidence indicates that mitochondrial involvement in $[Ca^{2+}]_i$ oscillations must also be related to ROS formation by oxidative phosphorylation.

These results suggest that any Ca^{2+} mobilizing signal would become a direct stimulus for the metabolic enzymes in the mitochondria, leading to activation of metabolic pathways and respiratory chain through multiple Ca^{2+} -regulated enzymes of the mitochondrial matrix (for review see Ref. 6). Once reduced equivalents are increased the amount of O_2^- increases, "pushed" by a higher number of electrons flowing from complex I and II to complex III. Indeed, there is correlation between metabolic rate and ROS generation (62). Therefore, mitochondria are suited to respond to cellular stimulation with an increase in ROS production. Agonist-induced mitochondria activation serves not only to supply the immediate energy required to extrude Ca^{2+} from cytosol (via SERCA and PMCA pumps) but also facilitates Ca^{2+} signals sensitizing the intracellular Ca^{2+} release channels.

The spatial arrangement of mitochondria with respect to Ca^{2+} release channels (IP₃R and RyR), which forms a privileged pathway for calcium transfer between reticulum and mitochondria (30, 67, 69), would allow fast diffusion of ROS from mitochondria to Ca^{2+} channels. In fact, the close apposition between mitochondria and the locus of endoplasmic reticulum rich in IP₃R and RyR is interpreted as a strategy for direct manipulation of the environment surrounding the Ca^{2+} channels' mouth. This has led to the concept of "intracellular synapse" (16). However, the mitochondrial buffer mechanism cannot explain some experimental results such as the inability of cytosolic buffers to mimic the effect of mitochondrial uptake (27).

The evidence presented above is not a negation of the "conventional" role of mitochondrial Ca^{2+} uptake effect on Ca^{2+} signals. In any case, the model presented here is compatible with the mitochondrial modulation of $[Ca^{2+}]_i$ around endoplasmic reticulum Ca^{2+} channels. Following the concept of "quasi-synaptic" transmission between mitochondria and reticulum coined by Hajnozcky's group (16), controlled amounts of mitochondrial oxidants could behave as a "neuromodulator" of the local effect of the released Ca^{2+} . In the same line, the multiple antioxidant systems present in healthy cells would terminate this "synaptic" signal, avoiding uncontrolled amplification by Ca^{2+} /ROS positive feedback. This is due not only to antioxidant barriers (SOD, catalase, etc.) but also to the high Ca^{2+} buffering capacity of the mitochondrial matrix, which limits the amplitude of mitochondrial Ca^{2+} transients.

An additional possibility is that endogenous ROS could recruit the cADP ribose pathway, because several groups have shown that cADP ribose metabolism can be modulated by redox mechanisms (58). This raises the possibility that once oscillations are started by IP₃ cADP ribose may cooperate to ensure their stabilization, given the sensitizing effect of this messenger on calcium signals evoked by agonists (11).

If mitochondrial ROS are confirmed as a modulatory factor in the operation of Ca²⁺ release channels under cellular stimulation, their presence must be taken into account in the development of models for Ca²⁺ signals and mitochondrial biology. Two aspects should be considered when modeling ROS and [Ca²⁺]_i signals. The most obvious is to incorporate ROS effects to [Ca²⁺]_i signaling models. Mitochondria have already been included in some calcium dynamic models (see, e.g., Ref. 50). In addition, Cortassa et al. (15) have described a detailed model of oscillations for mitochondrial O₂^{-•} production, inner membrane potential, and NADH, reproducing and predicting experimental observations performed in mitochondria from living cardiomyocytes. This model is based on a previous model from the same group describing the effect of mitochondrial calcium dynamics on energy metabolism (14). Therefore, incorporation of some aspects of this model for cytosolic [Ca²⁺]_i oscillations should give new insights into the mechanisms underlying this process and could predict the limits of the modulatory action of ROS on [Ca²⁺]_i signals.

Another issue to be considered is the effect of calcium ions on mitochondrial parameters, especially ROS production. This point has raised some uncertainty, given that mechanistic predictions indicate that [Ca²⁺]_m increases would not favor ROS production (15, 55) because they partially collapse mitochondrial potential (10, 28, 78, 87). Thus recent work has detected no effect or a slight decrease in ROS generation in response to [Ca²⁺]_m increases in isolated mitochondria (78), and Aon et al. (2, 3) report that ROS production is independent of Ca²⁺ uptake.

However, the same reports show Ca²⁺-induced ROS formation under partial depolarization, when this process is independent from mitochondrial potential (78), and calcium can enhance ROS generation in the presence of some degree of mitochondrial inhibition (6). In fact, there is a clear mechanistic link between [Ca²⁺]_m increases and ROS generation. An increase in [Ca²⁺]_m in metabolic situations where the NADH/FADH₂ production controls the electron transport rate enhances the respiratory flux and ROS production. The latter could be minimized in state 4 conditions by the Ca²⁺-induced dissipation of mitochondrial potential.

Mathematical models can help us to uncover the complex relationship between mitochondrial parameters and calcium homeostasis. In addition, this approach is necessary to understand how subtle and complex changes underlie the shift from normal to pathological states, for example, when the simultaneous presence of mitochondrial Ca²⁺ increases and an otherwise nonpathological factor leads to mitochondrial dysfunction (6). This approach would support the development of new therapies or strategies. This is extremely important in cellular models under energetic stress such as cardiac and brain tissues.

GRANTS

This work was supported by Ministerio de Educacion y Ciencia (BFU 2004-0637) and Junta de Extremadura (2PR03A020).

REFERENCES

1. Abramson JJ, Zable AC, Favero TG, and Salama G. Thimerosal interacts with the Ca²⁺ release channel ryanodine receptor from skeletal muscle sarcoplasmic reticulum. *J Biol Chem* 270: 29644–29647, 1995.
2. Aon MA, Cortassa S, Marban E, and O'Rourke B. Synchronized whole cell oscillations in mitochondrial metabolism triggered by a local release of reactive oxygen species in cardiac myocytes. *J Biol Chem* 278: 44735–44744, 2003.
3. Aon MA, Cortassa S, and O'Rourke B. Percolation and criticality in a mitochondrial network. *Proc Natl Acad Sci USA* 101: 4447–4452, 2004.
4. Beutner G, Sharma VK, Lin L, Ryu SY, Dirksen RT, and Sheu SS. Type 1 ryanodine receptor in cardiac mitochondria: transducer of excitation-metabolism coupling. *Biochim Biophys Acta* 1717: 1–10, 2005.
5. Bootman MD, Taylor CW, and Berridge MJ. The thiol reagent, thimerosal, evokes Ca²⁺ spikes in HeLa cells by sensitizing the inositol 1,4,5-trisphosphate receptor. *J Biol Chem* 267: 25113–25119, 1992.
6. Brookes PS, Yoon Y, Robotham JL, Anders MW, and Sheu SS. Calcium, ATP, and ROS: a mitochondrial love-hate triangle. *Am J Physiol Cell Physiol* 287: C817–C833, 2004.
7. Bultynck G, Szlufcik K, Kasri NN, Assefa Z, Callewaert G, Missiaen L, Parys JB, and De Smedt SH. Thimerosal stimulates Ca²⁺ flux through inositol 1,4,5-trisphosphate receptor type 1, but not type 3, via modulation of an isoform-specific Ca²⁺-dependent intramolecular interaction. *Biochem J* 381: 87–96, 2004.
8. Camello C, Camello PJ, Pariente JA, and Salido GM. Effects of antioxidants on calcium signal induced by cholecystokinin in mouse pancreatic acinar cells. *J Physiol Biochem* 56: 173–180, 2000.
9. Camello-Almaraz C, Salido GM, Pariente JA, and Camello PJ. Role of mitochondria in Ca²⁺ oscillations and shape of Ca²⁺ signals in pancreatic acinar cells. *Biochem Pharmacol* 63: 283–292, 2002.
10. Camello-Almaraz MC, Pozo MJ, Murphy MP, and Camello PJ. Mitochondrial production of oxidants is necessary for physiological calcium oscillations. *J Cell Physiol* 206: 487–494, 2006.
11. Cancela JM, Van CF, Galione A, Tepikin AV, and Petersen OH. Transformation of local Ca²⁺ spikes to global Ca²⁺ transients: the combinatorial roles of multiple Ca²⁺ releasing messengers. *EMBO J* 21: 909–919, 2002.
12. Chen Q, Vazquez EJ, Moghaddas S, Hoppel CL, and Lesnfsky EJ. Production of reactive oxygen species by mitochondria: central role of complex III. *J Biol Chem* 278: 36027–36031, 2003.
13. Coelho SM, Taylor AR, Ryan KP, Sousa-Pinto I, Brown MT, and Brownlee C. Spatiotemporal patterning of reactive oxygen production and Ca²⁺ wave propagation in *Fucus* rhizoid cells. *Plant Cell* 14: 2369–2381, 2002.
14. Cortassa S, Aon MA, Marban E, Winslow RL, and O'Rourke B. An integrated model of cardiac mitochondrial energy metabolism and calcium dynamics. *Biophys J* 84: 2734–2755, 2003.
15. Cortassa S, Aon MA, Winslow RL, and O'Rourke B. A mitochondrial oscillator dependent on reactive oxygen species. *Biophys J* 87: 2060–2073, 2004.
16. Csordas G, Thomas AP, and Hajnoczky G. Quasi-synaptic calcium signal transmission between endoplasmic reticulum and mitochondria. *EMBO J* 18: 96–108, 1999.
17. Dawson TL, Gores GJ, Nieminen AL, Herman B, and Lemasters JJ. Mitochondria as a source of reactive oxygen species during reductive stress in rat hepatocytes. *Am J Physiol Cell Physiol* 264: C961–C967, 1993.
18. Droge W. Free radicals in the physiological control of cell function. *Physiol Rev* 82: 47–95, 2002.
19. Droge W. Oxidative stress and aging. *Adv Exp Med Biol* 543: 191–200, 2003.
20. Duchon MR. Mitochondria and calcium: from cell signalling to cell death. *J Physiol* 529: 57–68, 2000.
21. Felty Q and Roy D. Estrogen, mitochondria, and growth of cancer and non-cancer cells. *J Carcinog* 4: 1, 2005.
22. Felty Q, Xiong WC, Sun D, Sarkar S, Singh KP, Parkash J, and Roy D. Estrogen-induced mitochondrial reactive oxygen species as signal-transducing messengers. *Biochemistry* 44: 6900–6909, 2005.
23. Feng W, Liu G, Allen PD, and Pessah IN. Transmembrane redox sensor of ryanodine receptor complex. *J Biol Chem* 275: 35902–35907, 2000.
24. Fink BD, Reszka KJ, Herlein JA, Mathahs MM, and Sivitz WI. Respiratory uncoupling by UCP1 and UCP2 and superoxide generation in

- endothelial cell mitochondria. *Am J Physiol Endocrinol Metab* 288: E71–E79, 2005.
25. Foreman MA, Smith J, and Publicover SJ. Characterisation of serum-induced intracellular Ca^{2+} oscillations in primary bone marrow stromal cells. *J Cell Physiol* 206: 664–671, 2006.
 26. Fusi F, Saponara S, Gagov H, and Sgaragli G. 2,5-Di-*t*-butyl-1,4-benzohydroquinone (BHQ) inhibits vascular L-type Ca^{2+} channel via superoxide anion generation. *Br J Pharmacol* 133: 988–996, 2001.
 27. Glitsch MD, Bakowski D, and Parekh AB. Store-operated Ca^{2+} entry depends on mitochondrial Ca^{2+} uptake. *EMBO J* 21: 6744–6754, 2002.
 28. Gonzalez A, Granados MP, Salido GM, and Pariente JA. Changes in mitochondrial activity evoked by cholecystokinin in isolated mouse pancreatic acinar cells. *Cell Signal* 15: 1039–1048, 2003.
 29. Gunter TE, Buntinas L, Sparagna G, Eliseev R, and Gunter K. Mitochondrial calcium transport: mechanisms and functions. *Cell Calcium* 28: 285–296, 2000.
 30. Hajnoczky G, Csordas G, Madesh M, and Pacher P. The machinery of local Ca^{2+} signalling between sarco-endoplasmic reticulum and mitochondria. *J Physiol* 529: 69–81, 2000.
 31. Hajnoczky G, Robb-Gaspers LD, Seitz MB, and Thomas AP. Decoding of cytosolic calcium oscillations in the mitochondria. *Cell* 82: 415–424, 1995.
 32. Hoth M, Fanger CM, and Lewis RS. Mitochondrial regulation of store-operated calcium signaling in T lymphocytes. *J Cell Biol* 137: 633–648, 1997.
 33. Hsieh TJ, Zhang SL, Filep JG, Tang SS, Ingelfinger JR, and Chan JS. High glucose stimulates angiotensinogen gene expression via reactive oxygen species generation in rat kidney proximal tubular cells. *Endocrinology* 143: 2975–2985, 2002.
 34. Hu Q, Yu ZX, Ferrans VJ, Takeda K, Irani K, and Ziegelstein RC. Critical role of NADPH oxidase-derived reactive oxygen species in generating Ca^{2+} oscillations in human aortic endothelial cells stimulated by histamine. *J Biol Chem* 277: 32546–32551, 2002.
 35. Huschenbett J, Zaidi A, and Michaelis ML. Sensitivity of the synaptic membrane Na^+/Ca^{2+} exchanger and the expressed NCX1 isoform to reactive oxygen species. *Biochim Biophys Acta* 1374: 34–46, 1998.
 36. Huser J and Blatter LA. Fluctuations in mitochondrial membrane potential caused by repetitive gating of the permeability transition pore. *Biochem J* 343: 311–317, 1999.
 37. Isaeva EV, Shkryl VM, and Shirokova N. Mitochondrial redox state and Ca^{2+} sparks in permeabilized mammalian skeletal muscle. *J Physiol* 565: 855–872, 2005.
 38. Izeradjene K, Douglas L, Tillman DM, Delaney AB, and Houghton JA. Reactive oxygen species regulate caspase activation in tumor necrosis factor-related apoptosis-inducing ligand-resistant human colon carcinoma cell lines. *Cancer Res* 65: 7436–7445, 2005.
 39. Jaggard JH, Porter VA, Lederer WJ, and Nelson MT. Calcium sparks in smooth muscle. *Am J Physiol Cell Physiol* 278: C235–C256, 2000.
 40. Kaftan EJ, Xu T, Abercrombie RF, and Hille B. Mitochondria shape hormonally induced cytoplasmic calcium oscillations and modulate exocytosis. *J Biol Chem* 275: 25465–25470, 2000.
 41. Kaplan P, Babusikova E, Lehotsky J, and Dobrota D. Free radical-induced protein modification and inhibition of Ca^{2+} -ATPase of cardiac sarcoplasmic reticulum. *Mol Cell Biochem* 248: 41–47, 2003.
 42. Kimura S, Zhang GX, Nishiyama A, Shokoji T, Yao L, Fan YY, Rahman M, and Abe Y. Mitochondria-derived reactive oxygen species and vascular MAP kinases: comparison of angiotensin II and diazoxide. *Hypertension* 45: 438–444, 2005.
 43. Kimura S, Zhang GX, Nishiyama A, Shokoji T, Yao L, Fan YY, Rahman M, Suzuki T, Maeta H, and Abe Y. Role of NAD(P)H oxidase- and mitochondria-derived reactive oxygen species in cardioprotection of ischemic reperfusion injury by angiotensin II. *Hypertension* 45: 860–866, 2005.
 44. Kirichok Y, Krapivinsky G, and Clapham DE. The mitochondrial calcium uniporter is a highly selective ion channel. *Nature* 427: 360–364, 2004.
 45. Kolisek M, Beck A, Fleig A, and Penner R. Cyclic ADP-ribose and hydrogen peroxide synergize with ADP-ribose in the activation of TRPM2 channels. *Mol Cell* 18: 61–69, 2005.
 46. Kourie JI. Interaction of reactive oxygen species with ion transport mechanisms. *Am J Physiol Cell Physiol* 275: C1–C24, 1998.
 47. Kushnareva Y, Murphy AN, and Andreyev A. Complex I-mediated reactive oxygen species generation: modulation by cytochrome *c* and NAD(P)⁺ oxidation-reduction state. *Biochem J* 368: 545–553, 2002.
 48. Lajas AI, Sierra V, Camello PJ, Salido GM, and Pariente JA. Vanadate inhibits the calcium extrusion in rat pancreatic acinar cells. *Cell Signal* 13: 451–456, 2001.
 49. Lehotsky J, Kaplan P, Matejovicova M, Murin R, Racay P, and Raeymaekers L. Ion transport systems as targets of free radicals during ischemia reperfusion injury. *Gen Physiol Biophys* 21: 31–37, 2002.
 50. Marhl M, Schuster S, and Brumen M. Mitochondria as an important factor in the maintenance of constant amplitudes of cytosolic calcium oscillations. *Biophys Chem* 2: 125–132, 1998.
 51. Missiaen L, Taylor CW, and Berridge MJ. Spontaneous calcium release from inositol trisphosphate-sensitive calcium stores. *Nature* 352: 241–244, 1991.
 52. Missiaen L, Taylor CW, and Berridge MJ. Luminal Ca^{2+} promoting spontaneous Ca^{2+} release from inositol trisphosphate-sensitive stores in rat hepatocytes. *J Physiol* 455: 623–640, 1992.
 53. Mittal CK and Murad F. Activation of guanylate cyclase by superoxide dismutase and hydroxyl radical: a physiological regulator of guanosine 3',5'-monophosphate formation. *Proc Natl Acad Sci USA* 74: 4360–4364, 1977.
 54. Montero M, Alonso MT, Albillos A, Garcia-Sancho J, and Alvarez J. Mitochondrial Ca^{2+} -induced Ca^{2+} release mediated by the Ca^{2+} uniporter. *Mol Biol Cell* 12: 63–71, 2001.
 55. Nicholls DG. Mitochondria and calcium signaling. *Cell Calcium* 38: 311–317, 2005.
 56. Nicholls DG and Budd SL. Mitochondria and neuronal survival. *Physiol Rev* 80: 315–360, 2000.
 57. Nicholls DG and Ward MW. Mitochondrial membrane potential and neuronal glutamate excitotoxicity: mortality and millivolts. *Trends Neurosci* 23: 166–174, 2000.
 58. Okabe E, Tsujimoto Y, and Kobayashi Y. Calmodulin and cyclic ADP-ribose interaction in Ca^{2+} signaling related to cardiac sarcoplasmic reticulum: superoxide anion radical-triggered Ca^{2+} release. *Antioxid Redox Signal* 2: 47–54, 2000.
 59. O'Reilly CM, Fogarty KE, Drummond RM, Tuft RA, and Walsh JV Jr. Quantitative analysis of spontaneous mitochondrial depolarizations. *Biophys J* 85: 3350–3357, 2003.
 60. Park MK, Ashby MC, Erdemli G, Petersen OH, and Tepikin AV. Perinuclear, perigranular and sub-plasmalemmal mitochondria have distinct functions in the regulation of cellular calcium transport. *EMBO J* 20: 1863–1874, 2001.
 61. Parkash J, Felty Q, and Roy D. Estrogen exerts a spatial and temporal influence on reactive oxygen species generation that precedes calcium uptake in high-capacity mitochondria: implications for rapid nongenomic signaling of cell growth. *Biochemistry* 45: 2872–2881, 2006.
 62. Perez-Campo R, Lopez-Torres M, Cadenas S, Rojas C, and Barja G. The rate of free radical production as a determinant of the rate of aging: evidence from the comparative approach. *J Comp Physiol [B]* 168: 149–158, 1998.
 63. Petersen CC, Toescu EC, and Petersen OH. Different patterns of receptor-activated cytoplasmic Ca^{2+} oscillations in single pancreatic acinar cells: dependence on receptor type, agonist concentration and intracellular Ca^{2+} buffering. *EMBO J* 10: 527–533, 1991.
 64. Pozo MJ, Perez GJ, Nelson MT, and Mawe GM. Ca^{2+} sparks and BK currents in gallbladder myocytes: role in CCK-induced response. *Am J Physiol Gastrointest Liver Physiol* 282: G165–G174, 2002.
 65. Pueyo ME, Gonzalez W, Nicoletti A, Savoie F, Arnal JF, and Michel JB. Angiotensin II stimulates endothelial vascular cell adhesion molecule-1 via nuclear factor- κ B activation induced by intracellular oxidative stress. *Arterioscler Thromb Vasc Biol* 20: 645–651, 2000.
 66. Rizzuto R, Bernardi P, and Pozzan T. Mitochondria as all-round players of the calcium game. *J Physiol* 529: 37–47, 2000.
 67. Rizzuto R, Brini M, Murgia M, and Pozzan T. Microdomains with high Ca^{2+} close to IP_3 -sensitive channels that are sensed by neighboring mitochondria. *Science* 262: 744–747, 1993.
 68. Rizzuto R, Duchon MR, and Pozzan T. Flirting in little space: the ER/mitochondria Ca^{2+} liaison. *Sci STKE* 2004: re1, 2004.
 69. Rizzuto R, Pinton P, Carrington W, Fay FS, Fogarty KE, Lifshitz LM, Tuft RA, and Pozzan T. Close contacts with the endoplasmic reticulum as determinants of mitochondrial Ca^{2+} responses. *Science* 280: 1763–1766, 1998.
 70. Rizzuto R, Simpson AW, Brini M, and Pozzan T. Rapid changes of mitochondrial Ca^{2+} revealed by specifically targeted recombinant aequorin. *Nature* 358: 325–327, 1992.

71. **Rutter GA, Burnett P, Rizzuto R, Brini M, Murgia M, Pozzan T, Tavare JM, and Denton RM.** Subcellular imaging of intramitochondrial Ca^{2+} with recombinant targeted aequorin: significance for the regulation of pyruvate dehydrogenase activity. *Proc Natl Acad Sci USA* 93: 5489–5494, 1996.
72. **Scanlon JM, Brocard JB, Stout AK, and Reynolds IJ.** Pharmacological investigation of mitochondrial Ca^{2+} transport in central neurons: studies with CGP-37157, an inhibitor of the mitochondrial Na^+ - Ca^{2+} exchanger. *Cell Calcium* 28: 317–327, 2000.
73. **Simpson PB.** The local control of cytosolic Ca^{2+} as a propagator of CNS communication—integration of mitochondrial transport mechanisms and cellular responses. *J Bioenerg Biomembr* 32: 5–13, 2000.
74. **Simpson PB and Russell JT.** Mitochondria support inositol 1,4,5-trisphosphate-mediated Ca^{2+} waves in cultured oligodendrocytes. *J Biol Chem* 271: 33493–33501, 1996.
75. **Sparagna GC, Gunter KK, Sheu SS, and Gunter TE.** Mitochondrial calcium uptake from physiological-type pulses of calcium. A description of the rapid uptake mode. *J Biol Chem* 270: 27510–27515, 1995.
76. **Spat A and Pitter JG.** The effect of cytoplasmic Ca^{2+} signal on the redox state of mitochondrial pyridine nucleotides. *Mol Cell Endocrinol* 215: 115–118, 2004.
77. **Starkov AA and Fiskum G.** Regulation of brain mitochondrial H_2O_2 production by membrane potential and NAD(P)H redox state. *J Neurochem* 86: 1101–1107, 2003.
78. **Starkov AA, Polster BM, and Fiskum G.** Regulation of hydrogen peroxide production by brain mitochondria by calcium and Bax. *J Neurochem* 83: 220–228, 2002.
79. **Straub SV, Giovannucci DR, and Yule DI.** Calcium wave propagation in pancreatic acinar cells: functional interaction of inositol 1,4,5-trisphosphate receptors, ryanodine receptors, and mitochondria. *J Gen Physiol* 116: 547–560, 2000.
80. **Suzuki YJ and Ford GD.** Redox regulation of signal transduction in cardiac and smooth muscle. *J Mol Cell Cardiol* 31: 345–353, 1999.
81. **Thorn P, Brady P, Llopis J, Gallacher DV, and Petersen OH.** Cytosolic Ca^{2+} spikes evoked by the thiol reagent thimerosal in both intact and internally perfused single pancreatic acinar cells. *Pflügers Arch* 422: 173–178, 1992.
82. **Tinel H, Cancela JM, Mogami H, Gerasimenko JV, Gerasimenko OV, Tepikin AV, and Petersen OH.** Active mitochondria surrounding the pancreatic acinar granule prevent spreading of inositol trisphosphate-evoked local cytosolic Ca^{2+} signals. *EMBO J* 18: 4999–5008, 1999.
83. **Turrens JF.** Mitochondrial formation of reactive oxygen species. *J Physiol* 552: 335–344, 2003.
84. **Turrens JF, Alexandre A, and Lehninger AL.** Ubisemiquinone is the electron donor for superoxide formation by complex III of heart mitochondria. *Arch Biochem Biophys* 237: 408–414, 1985.
85. **Vergun O and Reynolds IJ.** Fluctuations in mitochondrial membrane potential in single isolated brain mitochondria: modulation by adenine nucleotides and Ca^{2+} . *Biophys J* 87: 3585–3593, 2004.
86. **Voronina S, Sukhomlin T, Johnson PR, Erdemli G, Petersen OH, and Tepikin A.** Correlation of NADH and Ca^{2+} signals in mouse pancreatic acinar cells. *J Physiol* 539: 41–52, 2002.
87. **Voronina SG, Barrow SL, Gerasimenko OV, Petersen OH, and Tepikin AV.** Effects of secretagogues and bile acids on mitochondrial membrane potential of pancreatic acinar cells: comparison of different modes of evaluating $\Delta\Psi_m$. *J Biol Chem* 279: 27327–27338, 2004.
88. **Waypa GB, Marks JD, Mack MM, Boriboun C, Mungai PT, and Schumacker PT.** Mitochondrial reactive oxygen species trigger calcium increases during hypoxia in pulmonary arterial myocytes. *Circ Res* 91: 719–726, 2002.
89. **Waypa GB and Schumacker PT.** Hypoxic pulmonary vasoconstriction: redox events in oxygen sensing. *J Appl Physiol* 98: 404–414, 2005.
90. **Wu J, Takeo T, Kamimura N, Wada J, Suga S, Hoshina Y, and Wakui M.** Thimerosal modulates the agonist-specific cytosolic Ca^{2+} oscillatory patterns in single pancreatic acinar cells of mouse. *FEBS Lett* 390: 149–152, 1996.
91. **Xi Q, Cheranov SY, and Jaggar JH.** Mitochondria-derived reactive oxygen species dilate cerebral arteries by activating Ca^{2+} sparks. *Circ Res* 97: 354–362, 2005.
92. **Zaidi A and Michaelis ML.** Effects of reactive oxygen species on brain synaptic plasma membrane Ca^{2+} -ATPase. *Free Radic Biol Med* 27: 810–821, 1999.
93. **Zmijewski JW, Landar A, Watanabe N, Dickinson DA, Noguchi N, and Darley-Usmar VM.** Cell signalling by oxidized lipids and the role of reactive oxygen species in the endothelium. *Biochem Soc Trans* 33: 1385–1389, 2005.